Malaysian Cat Gecko Care Guide

Aeluroscalabotes felinus

Create a cool, heavily covered forest-understory habitat with gentle climbing surfaces, consistent hydration and good airflow. Locality and individual sensitivity matter; unverified claims that this species requires mineral-free drinking water should not guide care.

Species Profile

Common name: Malaysian Cat Gecko

Scientific name: *Aeluroscalabotes felinus*

Adult size: About 4-7 in (10-18 cm) total length; females larger

Average lifespan: 8-10 years

Native range: Southeast Asia, including Peninsular Malaysia, Borneo and nearby regions

Habitat and activity: Nocturnal; humid forest understory, low vegetation and leaf litter

Diet: Small varied cultured insects

Care difficulty: Advanced

Shopping List

- Secure enclosure at least 18 × 18 × 24 in (46 × 46 × 61 cm; L × W × H) for one adult; see Housing for exceptions and group arrangements.
- Digital thermometers for different zones; a suitable hygrometer; timer for a consistent day/night cycle.
- Species-appropriate substrate and multiple stable hides or covered retreats.
- Reliable supply of the staple foods described below, clean feeding tools and a transport container.
- Appropriate overhead heat source if required, controlled by a compatible thermostat; infrared thermometer for surfaces.
- Appropriate visible lighting and, where recommended below, a linear UVB fixture with measured shade and exposure zones.
- Fresh-water dish or cup; insect supplements when applicable; a scale appropriate to the animal's size.

Housing

Our recommended starting adult enclosure is 18 × 18 × 24 in (46 × 46 × 61 cm; L × W × H) for one animal. Larger usable space is encouraged. These dimensions are a husbandry recommendation, not a species-specific experimentally proven threshold.

Fill the height with firmly secured horizontal and diagonal branches, cork, foliage and sheltered paths. Provide several resting places at different heights and enough cover to travel between environmental zones without crossing a bare open box.

Juveniles may use a furnished adult enclosure when food intake, hydration and waste can be monitored. A temporary grow-out enclosure should still offer choice and cover. Quarantine newcomers separately with dedicated tools and a substrate that permits health monitoring; arrange veterinary assessment for unknown-origin animals.

Lighting & UVB

Provide roughly 10-12 hours of visible daylight followed by darkness. A measured UV Index (UVI) around 0.3-0.7 at the most exposed usable site, declining to near zero inside retreats, is our conservative starting recommendation. Low-level UVB is recommended when a usable gradient can be provided; nocturnal activity alone is not a reason to exclude light.

Measure exposure at the animal's back height with an appropriate UVI meter, such as a Solarmeter 6.5. Lamp percentage, wattage and a fixed distance alone do not establish exposure: reflectors, mesh, perch height, plant growth and lamp age change output. Keep glass/plastic out of the UV path, follow fixture safety instructions and provide complete shade. Replace or reassess lamps according to measured output and manufacturer guidance.

Heating & Temperature

- Warm retreat / local air: 76-80°F (24-27°C).
- Cool-zone air: 70-74°F (21-23°C).
- Nighttime air: 66-72°F (19-22°C) for healthy established animals.

Create the gradient on one side rather than heating every hide to the same temperature. Use a compatible thermostat on each heat source; a manual lamp dimmer does not replace thermostatic protection. Secure the probe and verify temperatures independently: probe thermometers measure air/retreat temperature, while an infrared thermometer measures exposed surfaces.

Use appropriately guarded overhead warmth when needed. Keep animals clear of hot fixtures, avoid heat rocks, and switch visible lights off at night. Night heat is needed only when the room falls below the appropriate range; use a thermostat-controlled lightless source if necessary.

Do not allow the whole enclosure to approach the warm-site temperature. Sustained mid-80s°F (about 29°C) air is a warning for these setups; heat-sensitive individuals may need intervention sooner. Keep a cooler refuge available and respond to prolonged heat avoidance.

Humidity & Ventilation

65-80% with temporary evening rises and airflow. Treat readings as a starting pattern rather than a number to chase regardless of substrate, airflow or animal behavior.

Measure humidity in a representative occupied zone, away from a freshly sprayed surface. Provide sheltered moisture choices while retaining airflow. Persistent condensation, sour odor, flooded soil or surfaces that never dry indicate a setup problem. Ventilation and hydration must be adjusted together rather than simply sealing all vents.

Water & Hydration

Provide a clean, appropriately shallow dish or accessible elevated water cup at all times. Change it daily and immediately when soiled; wash it regularly with dedicated tools. Potable water is generally suitable, with treatment where local water quality requires it.

For animals that drink droplets, mist foliage or suitable surfaces at the time they become active, then allow partial drying. Droplets supplement the dish; water chemistry claims should be based on evidence rather than a species' trade name.

Substrate

Because Malaysian cat geckos need high humidity, the substrate should hold moisture without becoming waterlogged. A good substrate mix is 50% organic topsoil and 50% coco fiber, with untreated sphagnum moss and clean leaf litter added on top. A drainage layer is strongly recommended, especially in bioactive setups or enclosures that are misted heavily.

The substrate should be damp in the lower layers, but the surface should not stay soaking wet. Leaf litter, cork pieces, and moss patches help create humid microclimates where the gecko can hide during the day.

Feces and urates should be removed as noticed. In non-bioactive enclosures, contaminated substrate should be scooped out and replaced, with a full substrate change every 1-3 months depending on cleanliness and moisture levels. In bioactive enclosures, maintain springtails/isopods, replace leaf litter as needed, and monitor for mold, sour smells, or saturated soil.

Food & Feeding

Malaysian cat geckos are strict insectivores, which means they eat insects and other small invertebrates. They should not be fed crested gecko diet, fruit puree, baby food, vegetables, or human food.

Good feeders include small crickets, small dubia roaches, small discoid roaches, isopods, black soldier fly larvae, silkworms, and small hornworms. Captive-raised snails may also be accepted by some individuals, but wild snails should not be used because of parasite and pesticide risk.

Feed at night or close to lights-out, when the gecko is naturally active.

How often Malaysian cat geckos need to eat depends on age:

Juveniles - 2-3 small insects, 4-5x/week

Adults - 2-3 appropriately sized insects, 2-3x/week

Prey should be no wider than the space between the gecko's eyes. Avoid leaving large numbers of crickets loose in the enclosure, as uneaten feeders can stress or bite the gecko.

Supplements

Lightly dust appropriate insect meals with calcium and use a balanced reptile multivitamin containing suitable vitamin A according to the product’s directions. Gut-loading and variety support nutrition but do not eliminate the need to balance calcium-poor prey.

Match D3 supplementation to the actual UVB exposure and to vitamins already included in other products. UVB does not automatically mean zero dietary D3, and lack of UVB does not justify unlimited D3. Follow one coherent program; do not combine multiple strong vitamin products or coat every feeder heavily.

Decor & Enrichment

Decorations play a major role in Malaysian cat gecko care. This is a shy, secretive species that feels safer when the enclosure is heavily covered. Good decor can include cork bark tubes, cork flats, thin branches, bamboo, vines, broad artificial leaves, live plants, moss patches, hanging hides, and leaf litter.

The enclosure should have cover at multiple levels, not just on the floor. The more secure the gecko feels, the more likely it is to eat well and move around naturally at night.

Handling & Behavior

Treat this animal primarily as a display species. For transfers, gently guide it into a cup, tube or transport container instead of chasing and grabbing. Keep doors and escape routes controlled before opening the enclosure.

Allow new animals to settle and eat reliably. Avoid restraint by the tail, repeated uncovering or forced interactions. Track normal activity for the individual so changes in feeding, posture or hiding can be recognized.

Cleaning & Routine Maintenance

Daily: check the animal's access to water/appropriate moisture, temperatures, visible condition and equipment. Remove feces, contaminated material, dead feeders and spoiled food promptly.

Regularly: clean dishes and feeding surfaces, check thermostats and probes, inspect decor stability and maintain ventilation. Bioactive systems still require waste checks, food removal, plant care and drainage maintenance; they are not self-cleaning.

Record appetite, activity, shed history and weight or body condition when safe. Arrange veterinary assessment for unexplained changes. Quarantine new animals with dedicated tools, and disinfect only with a suitable product used at the label dilution/contact time, then rinse and dry as required before return.

Cohabitation

Individual housing is the safest routine default. Shared housing can cause food competition, injury or reproductive pressure even without obvious fighting. Never assume a male/female pair or female group is compatible. Breeding introductions belong in a planned specialist program with separate backup enclosures and individual monitoring.

Shedding & Seasonal Changes

Maintain access to a clean humid retreat and inspect toes, tail tip and eyes after shedding when observation is possible. Do not peel retained skin or apply oils/adhesives. Persistent constriction, damaged tissue or repeated failed sheds warrants veterinary help.

Seasonal changes in appetite or activity can occur. Check temperatures, hydration and weight before treating prolonged refusal as normal. Deliberate brumation or breeding cycles require healthy animals and species-specific preparation; pet care does not require forcing a seasonal shutdown.

Common Health Concerns

- Poor body condition or feeding decline: check the full gradient, hydration and food quality; arrange assessment for persistent refusal accompanied by weight loss.
- Calcium/vitamin imbalance: weakness, tremors, a soft or misshapen jaw or abnormal limbs require veterinary assessment; extra high-dose supplements are not a substitute for diagnosis.
- Skin, shed or injury problems: repeated retained skin, wounds, swelling or abnormal discoloration deserve prompt attention.
- Respiratory or systemic illness: labored/open-mouth breathing at rest, marked lethargy, loss of balance, persistent diarrhea or rapid deterioration require an experienced exotic veterinarian.

- Unknown-origin or wild-caught animals may carry parasites. Fecal testing and veterinary-directed treatment are preferable to indiscriminate preventive medication.