Spearpoint Leaf-Tailed Gecko / Ebenau's Gecko Care Guide

Uroplatus ebenaui

Dense low vegetation, leaf-like cover, humid refuges and a safe thermal gradient matter more than a generic tropical-gecko formula. Verify the identification: several superficially similar leaf-tailed species have different requirements.

Species Profile

Common name: Spearpoint Leaf-Tailed Gecko / Ebenau's Gecko

Scientific name: *Uroplatus ebenaui*

Adult size: About 3-4 in (8-10 cm) total length

Average lifespan: 5-10 years

Native range: Northern and northwestern Madagascar and nearby islands

Habitat and activity: Nocturnal shrub/low-vegetation gecko; leaf-mimicking, shaded forest retreats

Diet: Small varied cultured insects

Care difficulty: Advanced

Shopping List

- Secure enclosure at least 18 × 18 × 24 in (46 × 46 × 61 cm; L × W × H) for one adult; see Housing for exceptions and group arrangements.
- Digital thermometers for different zones; a suitable hygrometer; timer for a consistent day/night cycle.
- Species-appropriate substrate and multiple stable hides or covered retreats.
- Reliable supply of the staple foods described below, clean feeding tools and a transport container.
- Appropriate overhead heat source if required, controlled by a compatible thermostat; infrared thermometer for surfaces.
- Appropriate visible lighting and, where recommended below, a linear UVB fixture with measured shade and exposure zones.
- Fresh-water dish or cup; insect supplements when applicable; a scale appropriate to the animal's size.

Housing

Our recommended starting adult enclosure is 18 × 18 × 24 in (46 × 46 × 61 cm; L × W × H) for one animal. Larger usable space is encouraged. These dimensions are a husbandry recommendation, not a species-specific experimentally proven threshold.

Fill the height with firmly secured horizontal and diagonal branches, cork, foliage and sheltered paths. Provide several resting places at different heights and enough cover to travel between environmental zones without crossing a bare open box.

Juveniles may use a furnished adult enclosure when food intake, hydration and waste can be monitored. A temporary grow-out enclosure should still offer choice and cover. Quarantine newcomers separately with dedicated tools and a substrate that permits health monitoring; arrange veterinary assessment for unknown-origin animals.

Lighting & UVB

Provide roughly 10-12 hours of visible daylight followed by darkness. A measured UV Index (UVI) around 0.3-0.7 at the most exposed usable site, declining to near zero inside retreats, is our conservative starting recommendation. Low-level UVB is recommended when a usable gradient can be provided; nocturnal activity alone is not a reason to exclude light.

Measure exposure at the animal's back height with an appropriate UVI meter, such as a Solarmeter 6.5. Lamp percentage, wattage and a fixed distance alone do not establish exposure: reflectors, mesh, perch height, plant growth and lamp age change output. Keep glass/plastic out of the UV path, follow fixture safety instructions and provide complete shade. Replace or reassess lamps according to measured output and manufacturer guidance.

Heating & Temperature

- Warm retreat / local air: 76-80°F (24-27°C).
- Cool-zone air: 70-74°F (21-23°C).
- Nighttime air: 66-72°F (19-22°C) for healthy established animals.

Create the gradient on one side rather than heating every hide to the same temperature. Use a compatible thermostat on each heat source; a manual lamp dimmer does not replace thermostatic protection. Secure the probe and verify temperatures independently: probe thermometers measure air/retreat temperature, while an infrared thermometer measures exposed surfaces.

Use appropriately guarded overhead warmth when needed. Keep animals clear of hot fixtures, avoid heat rocks, and switch visible lights off at night. Night heat is needed only when the room falls below the appropriate range; use a thermostat-controlled lightless source if necessary.

Do not allow the whole enclosure to approach the warm-site temperature. Sustained mid-80s°F (about 29°C) air is a warning for these setups; heat-sensitive individuals may need intervention sooner. Keep a cooler refuge available and respond to prolonged heat avoidance.

Humidity & Ventilation

Around 60% while drying and 80% after evening misting; keep humid retreats. Treat readings as a starting pattern rather than a number to chase regardless of substrate, airflow or animal behavior.

Measure humidity in a representative occupied zone, away from a freshly sprayed surface. Provide sheltered moisture choices while retaining airflow. Persistent condensation, sour odor, flooded soil or surfaces that never dry indicate a setup problem. Ventilation and hydration must be adjusted together rather than simply sealing all vents.

Water & Hydration

Provide a clean, appropriately shallow dish or accessible elevated water cup at all times. Change it daily and immediately when soiled; wash it regularly with dedicated tools. Potable water is generally suitable, with treatment where local water quality requires it.

For animals that drink droplets, mist foliage or suitable surfaces at the time they become active, then allow partial drying. Droplets supplement the dish; water chemistry claims should be based on evidence rather than a species' trade name.

Substrate

A moisture-holding tropical substrate works best. A good DIY mix is 50% organic topsoil and 50% coco fiber, with clean leaf litter and patches of untreated sphagnum moss on top. A drainage layer is strongly recommended because this species needs regular misting.

The substrate should help support humidity without becoming sour, swampy, or constantly saturated. Leaf litter is especially useful because it creates natural hiding areas, supports microfauna, and helps the surface dry more evenly.

Feces and urates should be removed when noticed. In a non-bioactive enclosure, contaminated substrate should be scooped out and replaced, with a full substrate change every 1-3 months depending on moisture and cleanliness. Bioactive setups need springtails, isopods, leaf litter replacement, and regular monitoring.

Food & Feeding

Spearpoint leaf-tailed geckos are strict insectivores, which means that they eat insects and should not be fed crested gecko diet, fruit, baby food, or vegetables. Feed at night or just before lights out.

Good feeder insects include small crickets, small roaches, black soldier fly larvae, silkworms, small hornworms, and other appropriately sized soft-bodied feeders. Prey should be small, usually no wider than the space between the gecko's eyes.

How often spearpoint leaf-tailed geckos need to eat depends on age:

Juveniles - small insects 4-5x/week

Adults - small insects 2-3x/week

Offer only what the gecko will eat in a short period, and remove uneaten crickets. Loose crickets can stress or injure a resting leaf-tailed gecko.

Supplements

Lightly dust appropriate insect meals with calcium and use a balanced reptile multivitamin containing suitable vitamin A according to the product’s directions. Gut-loading and variety support nutrition but do not eliminate the need to balance calcium-poor prey.

Match D3 supplementation to the actual UVB exposure and to vitamins already included in other products. UVB does not automatically mean zero dietary D3, and lack of UVB does not justify unlimited D3. Follow one coherent program; do not combine multiple strong vitamin products or coat every feeder heavily.

Decor & Enrichment

Decorations are not optional for spearpoint leaf-tailed geckos. The enclosure should be dense enough that the gecko can hide while still having room to move. Cork bark, thin branches, vines, dead leaves, broad foliage, live plants, and textured backgrounds all help this species behave naturally.

Provide both vertical and diagonal perches, but keep the layout focused on low to mid-level cover rather than only high branches. The gecko should be able to rest against bark, leaves, or cork in several areas with different humidity and temperature levels.

Handling & Behavior

Treat this animal primarily as a display species. For transfers, gently guide it into a cup, tube or transport container instead of chasing and grabbing. Keep doors and escape routes controlled before opening the enclosure.

Allow new animals to settle and eat reliably. Avoid restraint by the tail, repeated uncovering or forced interactions. Track normal activity for the individual so changes in feeding, posture or hiding can be recognized.

Cleaning & Routine Maintenance

Daily: check the animal's access to water/appropriate moisture, temperatures, visible condition and equipment. Remove feces, contaminated material, dead feeders and spoiled food promptly.

Regularly: clean dishes and feeding surfaces, check thermostats and probes, inspect decor stability and maintain ventilation. Bioactive systems still require waste checks, food removal, plant care and drainage maintenance; they are not self-cleaning.

Record appetite, activity, shed history and weight or body condition when safe. Arrange veterinary assessment for unexplained changes. Quarantine new animals with dedicated tools, and disinfect only with a suitable product used at the label dilution/contact time, then rinse and dry as required before return.

Cohabitation

Individual housing is the safest routine default. Shared housing can cause food competition, injury or reproductive pressure even without obvious fighting. Never assume a male/female pair or female group is compatible. Breeding introductions belong in a planned specialist program with separate backup enclosures and individual monitoring.

Shedding & Seasonal Changes

Maintain access to a clean humid retreat and inspect toes, tail tip and eyes after shedding when observation is possible. Do not peel retained skin or apply oils/adhesives. Persistent constriction, damaged tissue or repeated failed sheds warrants veterinary help.

Seasonal changes in appetite or activity can occur. Check temperatures, hydration and weight before treating prolonged refusal as normal. Deliberate brumation or breeding cycles require healthy animals and species-specific preparation; pet care does not require forcing a seasonal shutdown.

Common Health Concerns

- Poor body condition or feeding decline: check the full gradient, hydration and food quality; arrange assessment for persistent refusal accompanied by weight loss.
- Calcium/vitamin imbalance: weakness, tremors, a soft or misshapen jaw or abnormal limbs require veterinary assessment; extra high-dose supplements are not a substitute for diagnosis.
- Skin, shed or injury problems: repeated retained skin, wounds, swelling or abnormal discoloration deserve prompt attention.
- Respiratory or systemic illness: labored/open-mouth breathing at rest, marked lethargy, loss of balance, persistent diarrhea or rapid deterioration require an experienced exotic veterinarian.

- Unknown-origin or wild-caught animals may carry parasites. Fecal testing and veterinary-directed treatment are preferable to indiscriminate preventive medication.